



## Crop Recommendations for Cucumbers

### For Liquid Planters:

Apply Vitazyme with liquid starter fertilizer or water either directly in the furrow or 2x2 at the rate of 1 liter/hectare (13-16 oz/acre).

For small plots spray a 1% solution directly on the seeds in furrow before the row is closed at the rate of 1 liter/hectare (13-16 oz/acre).

### Dry Planters:

Apply Vitazyme with herbicide either before or after planting (either before or shortly after emergence) at the rate of 1 liter/hectare (13-16 oz/acre).

Apply a second time at early blossom, approximately 30 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).

Apply a third time approximately 60 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).

### Drip Irrigation:

Apply Vitazyme either before or after planting (either before or shortly after emergence) at the rate of 1 liter/hectare (13-16 oz/acre).

Apply a second time at early blossom, approximately 30 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).

Apply a third time approximately 60 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).

For extended yield apply after first picking at the rate of 1 liter/hectare (13-16 oz/acre).

***Vitazyme can be tank mixed with all farm chemicals, including herbicides, insecticides, fungicides, and fertilizers.***

***Added benefit: when mixed with herbicide, Vitazyme will stimulate weed growth, thereby enhancing herbicide efficacy.***