



Application Rates for Plums

New Orchards

Use one of the following treatments:

- Root dip: Dip the bare roots in a 5% Vitazyme solution
or
- Root Drench: During or after planting, drench the root zone with a 1% solution
or
- Spray Vitazyme at 16oz/acre (1 liter/hectare) with first foliar or weed spray

After planting, apply Vitazyme at the rate of 16oz/acre (1 liter/hectare) at approximately 30 day intervals during the growing season, 4 applications total.

Non Bearing Blocks

Apply at the rate of 16oz/acre (1 liter/hectare) with herbicide or with first foliar applied spray, and again at approximately 30 day intervals.

Bearing Blocks

Apply .80 oz/10,000 cu. ft. of Tree Row Volume (13-33 oz/acre or 1-2 liters/hectare) of Vitazyme, depending on the size of the trees. When in doubt use 1-1.5 pints per acre (1-1.5 liters/hectare):

- To the soil at budbreak or foliar at early pink
- at petal fall
- at first cover spray, or 30 days after petal fall
- 0-30 days post harvest

Vitazyme can be tank mixed with all farm chemicals, including herbicides, insecticides, fungicides, and fertilizers.

Added benefit: when mixed with herbicide, Vitazyme will stimulate weed growth, thereby enhancing herbicide efficacy