



Crop Recommendations for Watermelons

For Liquid Planters:

- Apply Vitazyme with liquid starter fertilizer or water either directly in the furrow or 2x2 at the rate of 1 liter/hectare (13-16 oz/acre).
- For small plots spray a 1% solution directly on the seeds in furrow before the row is closed at the rate of 1 liter/hectare (13-16 oz/acre).

Dry Planters:

- Apply Vitazyme with herbicide either before or after planting (either before or shortly after emergence) at the rate of 1 liter/hectare (13-16 oz/acre).
- Apply a second time at early blossom, approximately 30 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).
- Apply a third time approximately 60 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).
- Drip Irrigation:
- Apply Vitazyme either before or after planting (either before or shortly after emergence) at the rate of 1 liter/hectare (13-16 oz/acre).
- Apply a second time at early blossom, approximately 30 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).
- Apply a third time approximately 60 days after emergence at the rate of 1 liter/hectare (13-16 oz/acre).

Vitazyme can be tank mixed with all farm chemicals, including herbicides, insecticides, fungicides, and fertilizers.

Added benefit: when mixed with herbicide, Vitazyme will stimulate weed growth, thereby enhancing herbicide efficacy.